



## ***Streptococcus mutans* Dental Carries among Patients Attending Jimma University Specialized Hospital, Ethiopia**

Demissew Shenkute, Tsegahun Asfaw

Department of Medical Laboratory Science, Debre Berhan University, Debre Berhan, Ethiopia

**\*Corresponding Author:** Tsegahun Asfaw, Department of Medical Laboratory Science, Debre Berhan

**Abstract:** Dental caries is an irreversible microbial disease of the calcified tissues of the teeth. *Streptococcus mutans* is a bacterial resident of the oral cavity and is considered to be the principal etiological agent of dental caries in humans. Therefore, the aim of this study was to determine the prevalence and risk factors associated with *streptococcus mutans* dental caries. A cross sectional study was conducted among patients who attended Jimma University specialized hospital dental clinic. Patients demographic and clinical information was collected by using pre-tested questionnaire. Dental plaques from all patients was picked up by forceps and suspended in phosphate-buffered saline for further *streptococcus mutans* identification. From a total of 115 study participants 56(48.7%) and 59(51.3%) were males and females respectively. The overall prevalence of *streptococcus mutans* was 79(68.7%) among patients with dental carries. Twenty-five (21.7%) participants had gum bleeding and 47 (40.9%) participants had previous tooth decay. In this study, *streptococcus mutans* is the common public health problem among dental carries patients. Taking soft drinks, oral debris and gingival index were risk factor for dental caries of *streptococcus mutans*.

**Keywords:** *Streptococcus mutans*, Dental Carries, Ethiopia

### **1. INTRODUCTION**

Dental plaque is an adherent deposit of bacteria and their products, which forms as a white greenish or even yellow film on all tooth surfaces. Dental plaque accumulates naturally at stagnant or retentive sites formed after one or two days with no oral hygiene (Nada HA *et al.*, 2008). Dental caries is the single most prevalent and costly infectious disease worldwide, affecting more than 90% of the population in the US. The development of dental cavities requires the colonization of the tooth surface by acid producing bacteria such as *Streptococcus mutans*, in conjunction with the frequent ingestion of a cariogenic high sucrose diet, the substrate for acid and glucan production by organisms. The elevated amounts of acid and glucans modulate the establishment of cariogenic organisms within tightly adherent biofilms known as dental plaque (Catalan MA *et al.*, 2011). Tooth surfaces colonized with *S. mutans* are at a higher risk for developing caries (Pannu P. *et al.*, 2013). Dental caries is an irreversible microbial disease of the calcified tissues of the teeth, which is characterized by demineralization of the inorganic portion and destruction of the organic substance of the tooth, which often leads to cavitation. Dental caries is a multifactorial disease, which is caused by host, agent, and environmental factors (Kt S. *et al.*, 2013). *Streptococcus mutans* are gram positive cocci bacteria. These facultative anaerobes are commonly found in the human oral cavity. It grows at temperatures of between 18-40°C. *Streptococcus mutans* are a cariogenic microorganism that breaks down sugar for energy and produces an acidic environment, which dematerializes the superficial structure of the tooth. The result of the conversion disintegrates the coating of the tooth then later dissolves the calcium molecule creating a hole (Whiley RA. and Beighton D. 2013; Jav ND. *et al.*, 2013). It can be isolated from individuals either with or without a history of caries, but the development of dental caries is dependent on the bacterial load strain variation of *Streptococcus mutans* (Smith B *et al.*, 1998; Zhang L. *et al.*, 2009).

### **2. MATERIAL AND METHODS**

#### **2.1. Study Area and Period**

The study was conducted at Jimma University specialized hospital, which is located around 352 km Southwest of Addis Ababa. The study was conducted from March, 2017 to August, 2018.

## **2.2. Study Design**

A cross sectional study design was conducted.

## **2.3. Inclusion and Exclusion Criteria**

All patients attending dental clinic of Jimma University specialized hospital with clinical manifestation of dental caries and who are volunteer and able to give sample were included. Patients who are taking antibiotics in the last 30 days were excluded.

## **2.4. Sample Size and Sampling Technique**

The sampling technique was based on convenient sampling method. The sample size was all patients with clinical manifestation of dental caries who were visited Jimma University specialized hospital dental clinic and fulfill inclusion criteria.

## **2.5. Data Collection Methods**

Patients socio demographic and clinical information was collected by using pre-tested questionnaire. All study participants were examined by dental doctor. Dental plaques from all patients were picked up through forceps (probe) and transferred into 2 ml of sterile tube containing phosphate-buffered saline and processed immediately after collection in Jimma university microbiology laboratory (Teresa CC. *et al.*, 2007).

## **2.6. Isolation and Identification of *S. mutans***

The dental plaque suspension was vortexed for 30s for homogenization. Then a volume of 100µl was spread onto Mitis-salivaris (MS-agar) by using sterile cotton swab and incubated in 5% CO<sub>2</sub> for 48 hours at 37°C. A colony count of more than 250 (10<sup>4</sup> cells/ml) was considered as positive samples. Small colony was sub-cultured on blood-agar plates for further identification. A gram positive cocci, alpha hemolytic on blood agar, catalase negative, mannitol and lactose fermenters was considered as *Streptococcus mutans* (Nada HA. *et al.*, 2008; Cockerill FR. *et al.*, 2011).

## **2.7. Data Entry and Analysis**

Data was edited, cleaned and checked for its completeness and entered to SPSS version 21 for analysis. Patients' Socio-demographic and clinical characteristics were described by using descriptive statistics. Bivariate and multivariate logistic regression analysis were done and P-value less than 0.05 was taken as statistically significant.

## **2.8. Ethical Consideration**

Ethical clearance was obtained from Jimma University Ethical Review Board. Letter of permission was secured from Jimma University specialized hospital management. Written informed consent and assent was obtained from the study participants. Any information concerning the patients was kept confidential and patients with positive result were communicated for further management.

## **2.9. Quality Control**

To assure the quality of the data, standard operating procedure was followed during media preparation and other laboratory procedures. Sterility check was performed to avoid the possibility of contamination. All reagents were checked for their expiry date and prepared according to the manufacturer's instruction.

## **3. RESULT**

Among 115 study participants, the overall prevalence of *Streptococcus mutans* dental caries was 79 (68.7%).

### **3.1. Socio-demographic Characteristics of the of the Participants**

Among 115 participants, 56(48.7%) were males and 59(51.3%) were females. The mean age of the participants was 32.67 years with standard deviation (SD =1.48). Among participants 64 (55.7%) and 51(44.3%) of the participants were come from urban and rural area respectively. Regarding occupation 17(14.8%) of them were housewife followed by farmer 13(11.3%). Majority of the respondents 34(49.3%) were completed grade 12 (Table 1).

**Table1.** socio-demographic characteristics of patients attending dental clinic of Jimma University Specialized Hospital from March, 2017 to August, 2018

Socio- demographic Characteristics		Culture result for <i>Streptococcus mutans</i>		Total NO (%)
		Positive NO (%)	Negative NO (%)	
Sex	Male	43(37.4)	13(11.3)	56(48.7)
	Female	36(31.3)	23(20.0)	59(51.3)
Place of residence	Urban	41(35.7)	23(20.0)	64(55.7)
	Rural	38(33.0)	13(11.3)	51(44.3)
Occupation	Farmer	11(9.6)	2(1.7)	13(11.3)
	Merchant	6(5.2)	5(4.3)	11(9.6)
	Employed	10(8.7)	7(6.1)	17(14.8)
	Unemployed	3(2.6%)	1(0.9)	4(3.5)
	Housewife	11(9.6)	6(5.2)	17(14.8)
	Driver	1(0.9)	0(0)	1(0.9)
	Other	37(32.2)	15(13.0)	52(45.2)
Educational status	Illiterate	36(31.3)	10(8.7)	46(40.0)
	Read & write only	2(2.9)	3(4.3)	5(7.2)
	1-4 grade	5(7.2)	5(7.2)	10(14.5)
	5-8 grade	7(10.1)	1(1.4)	8(11.6)
	9-12 grade	7(10.1)	5(7.2)	12(17.4)
	Above 12	22(31.9)	12(17.4)	34(49.3)

### 3.2. Knowledge and Practice on Tooth Cleaning Habit of the Respondents

From 115 participants 73(63.5%) were cleaned their teeth and among these 73(63.5%), 40(54.8%) were used tooth stick and tooth brush respectively. From respondents who cleaned their teeth more than half of them were cleaned their teeth in sideways (horizontally) and also most of the participants were cleaned their tooth after meal and morning only. Very few were cleaned their teeth morning and before going to bed (Table 2).

**Table2.** Knowledge and practice on tooth cleaning habit of patients attending dental clinic of Jimma University Specialized Hospital from March, 2017 to April 2018

Knowledge and practice on oral health		Culture result for <i>Streptococcus mutans</i>		
		Positive NO (%)	Negative NO (%)	Total NO (%)
Habit of cleaning teeth	Yes	47(40.9)	26(22.6)	73(63.5)
	No	32(27.8)	10(8.7)	42(36.5)
Materials used to clean teeth	Tooth Stick	28(38.4)	12(16.4)	40(54.8)
	Charcoal	1(1.4)	2(2.7)	3(4.1)
	Tooth brush & rinse with water	9(12.3)	8(11.0)	17(23.3)
	Others	9(12.3)	4(5.5)	13(17.8)
Frequency of cleaning teeth	Once a day	22(30.1)	4(5.5)	26(35.6)
	After each meal	2(2.7)	3(4.1)	5(6.8)
	Before and after each meal	2(2.7)	3(4.1)	5(6.8)
	More than once a day	3(4.1)	4(5.5)	7(9.6)
	Irregularly	18(24.7)	12(16.4)	30(41.1)
Way of cleaning teeth	Top to bottom	6(8.2)	5(6.8)	11(15.1)
	Side way	23(31.5)	10(13.7)	33(45.2)
	Mixed	16(21.9)	11(15.1)	27(37.0)
	Circular	2(2.7)	0(0.0)	2(2.7)
Time of brushing teeth	Morning only	13(17.8)	10(13.7)	23(31.5)
	After meal	16(21.9)	8(11.0)	24(32.9)
	Before meal	3(4.1)	0(0.0)	3(4.1)
	Before going to bed	2(2.7)	0(0.0)	2(2.7)
	Irregular	11(15.1)	8(11.0)	19(26.1)

### 3.3. Associated Risk Factor for Dental Caries

Sixteen (13.9%) and 8(7.0%) of the study participants had a habit of drinking alcohol and smoking cigarette respectively. Twenty eight (24.3%) of the participants had a habit of chewing chat. Seventy

two (62.6%) of the participants had a habit of taking sweet in take or food. Twenty two (30.6%) of the participants were taking sweet intake irregularly. Eighty one (70.4%) of the participants had a habit of taking soft drink and significantly associated with *Streptococcusmutans* dental carries (p=0.01) (Table: 3).

**Table3.** Associated risk factor for dental caries in patients attending dental clinic of Jimma University Specialized Hospital from March, 2017 to April 2018

Associated risk factor		Culture result <i>Streptococcus mutans</i>			P-value
		Positive No (%)	Negative No (%)	Total No (%)	
Habit of drinking alcohol	Yes	13(11.3)	3(2.6)	16(13.9)	0.24
	No	66(57.4)	33(28.7)	99(86.1)	
Frequency of drinking alcohol	Always	0(0.0)	1(6.2)	1(6.2)	0.10
	Sometimes	6(37.5)	1(6.2)	7(43.8)	
	Rarely	7(43.8)	1(6.2)	8(50)	
Habit of smoking cigarette	Yes	8(7)	1(0.9)	9(7.8)	0.17
	No	71(61.7)	35(30.4)	106(92.2)	
Frequency of smoking cigarette	Regularly	5(55.6)	0(0.0)	5(55.6)	0.32
	Occasionally	3(33.3)	0(0.0)	3(33.3)	
	Rarely	0(0.0)	1(11.1)	1(11.1)	
Habit chewing chat	Yes	23(20)	5(4.3)	28(24.3)	0.08
	No	56(48.7)	31(27.0)	87(75.7)	
Frequency of chewing chat	Regularly	8(28.6)	1(3.6)	9(32.6)	0.40
	Occasionally	12(42.9)	2(7.1)	14(50.0)	
	Rarely	3(10.7)	2(7.1)	5(17.9)	
Habit of taking sweet intakes or food	Yes	46(40)	26(22.6)	72(62.6)	0.15
	No	33(28.7)	10(8.7)	43(37.4)	
Kind of sweet intake	Chocolate	0(0.0)	3(4.2)	3(4.2)	0.20
	Candy	1(1.4)	1(1.4)	2(2.8)	
	Sugar cane	6(8.3)	3(4.2)	9(12.5)	
	Sugared coffee	21(29.1)	8(11.1)	29(40.3)	
	Sugared tea	12(16.7)	9(12.5)	21(29.2)	
	Others	6(8.3)	2(2.8)	8(11.1)	
Frequency of taking sweet intake	Once a day	13(18.1)	3(4.2)	16(22.2)	0.15
	After each meal	8(11.1)	1(1.4)	9(12.5)	
	More than once a day	12(16.7)	10(13.9)	22(30.6)	
	Irregularly	12(16.7)	10(13.9)	22(30.6)	
	Every other day	1(1.4)	1(1.4)	2(2.8)	
	Once time per week	0(0.0)	1(1.4)	1(1.4)	
Habit of taking soft drinks	Yes	50(43.5)	31(27.0)	81(70.4)	0.01
	NO	29(25.2)	5(4.3)	34(9.6)	
frequency of taking soft drink	Regularly	3(3.7)	4(4.9)	7(8.6)	0.26
	Occasionally	28(34.6)	12(14.8)	40(49.4)	
	Rarely	19(23.5)	15(18.5)	34(42.0)	
presence xerostomia	Yes	4(3.5)	0(0.0)	4(3.5)	0.17
	No	75(65.2)	36(31.3)	111(100)	
Diabetes mellitus	Yes	15(13.0)	4(3.5)	19(16.5)	0.29
	No	64(55.7)	32(27.8)	96(83.5)	

### 3.4. Clinical Data on Oral Health

Among a total of 115 participants, 25(21.7%) had gem bleeding and 47 (40.9%) of the participants had previous tooth decay. Nearly more than half the respondents had oral debris covering not more than one third of the tooth surface. Only 42 (36%) were free from oral debris. Oral debris has statistically significance association with *Streptococcus mutans* dental caries (P = 0.022).

Most of the participants, 55(47.8%) had normal gingival index and 50(43.5%) had mild inflammation but only 1(0.9%) had sever inflammation. Gingival index has statistically significant association with *Streptococcus mutans* dental caries (P = 0.002).Fifty eight (50.4%) of the participant had no calculus index and41 (35.7%) had mild super gingival calculus. Only few 3(2.6%) had excessive supra and sub gingival calculus (Table 4).

**Table4.** Clinical finding in patients attending dental clinic of Jimma University Specialized Hospital from March, 2017 to April 2018

Clinical finding		Result			P-value
		Positive No (%)	Negative No (%)	Total No (%)	
Gem bleeding	Yes	18(15.7)	7(6.1)	25 (21.7)	0.69
	No	61(53.0)	29(25.2)	90(78.3)	
Previous tooth decay	Yes	35(30.4)	12(10.4)	47(40.9)	0.27
	No	44(38.3)	24(20.9)	68(59.1)	
Oral debris	No debris	22(19.1)	20(17.4)	42(16.5)	0.022
	Soft Debris covering not more than 1/3 of the tooth surface	47(40.9)	14(12.2)	64(53.0)	
	Soft debris covering more than 1/3 but not more than 2/3 of the exposed tooth surface	9(7.8)	1(0.9)	10(8.7)	
	soft debris covering more than two third of exposed tooth surface	1(0.9)	1(0.9)	2(1.8)	
Plaque index	no plaque	1(27.0)	20(17.4)	51(44.4)	0.23
	plaque adhering to the free gingival margin which cannot be seen by naked eye	39(33.9)	12(10.4)	51(44.3)	
	moderate accumulation of deposits on the gingival margin which can be seen with naked eye	9(7.8)	4(3.5)	13(11.3)	
gingival index	Normal	30(26.1)	25(21.7)	55(47.8)	0.002
	mild inflammation	40(34.8)	10(8.7)	50(43.5)	
	Moderate inflammation	9(7.8)	0(0)	9(7.8)	
	sever inflammation	0(0)	1(0.9)	1(0.9)	
calculus index	No calculus	36(31.3)	22(19.1)	58(50.4)	0.24
	Mild super gingival calculus	29(25.2)	12(10.4)	41(35.7)	
	Moderate supra & sub gingival calculus	11(9.6)	2(1.7)	13(11.3)	
	Excessive supra & sub gingival calculus	3(2.6)	0(0)	3(2.6)	

#### Class of Tooth Decay

From 47 tooth decay cases, 21 (44.7%) of them had class I followed by class II 17(36.2%) (Table 5).

**Table5.** Frequency of class of tooth decaying patients attending dental clinic of Jimma University Specialized Hospital with culture results for *Streptococcus mutans*

Class of tooth decay	Frequency No (%)
Class I	21 (44.7)
Class II	17 (36.2)
Class III	4 (8.5)
Class IV	2 (4.3)
Class V	3 (6.4)
Total	47 (100)

#### 4. DISCUSSION

In Ethiopia, there is scarcity of data on *Streptococcus mutans* dental caries. In this study *Streptococcus mutans* dental caries is the major public health problem among patients attending dental clinic of Jimma University specialized hospital. The prevalence of *Streptococcus mutans* dental caries was found to be 68.7% which is comparable with a study conducted in Brazil (68.5%) (Costa SM. *et al.*, 2012), China (67.5%) (Liu L. *et al.*, 2013) and Srilanka (68.8%) (Perera PJ. *et al.*, 2012). However, it was lower than study conducted in Qatar (85%) (Al-Darwish M. *et al.*, 2014). It was higher than study conducted in Gondar, Ethiopia (36.3%) (Ayele FA. *et al.*, 2013). The difference might be due to the difference in knowledge and practice of oral hygiene.

Habit of taking soft drink was found to be statistically significance association with *Streptococcus mutans* dental caries. This also supported by similar findings done in India and Zimbabwe respectively (Prasai Dixit L. *et al.*, 2013; Sukhabogi J. *et al.*, 2014). This also might be associated with acid production by cariogenic organism such as *Streptococcus mutans* that adhere to teeth as result of fermentation of soft drink. Later the enamel of tooth went to tooth decay.

In this study, the highest prevalence was seen in urban area 41(35.7%) than rural 38(33.0%). This finding was in agreements with a study done in Zimbabwe (Prasai Dixit L. *et al.*, 2013). The possible reason for this also might be due to the habit of urban population taking soft drink and food. Oral debris has statistically significant association with *Streptococcus mutans* dental caries. This finding also conforms other findings (Mafuvadze BT. *et al.*, 2013). Moreover; gingival index have significant association with *Streptococcus mutans* dental caries. This also might be a good indicator of poor oral hygiene practices (TS. Kumar *et al.*, 2013; Nobile CG. *et al.*, 2014). Because gingival index increase *Streptococcus mutans* colonization and in severe cases it involves loses of the enamel.

In this study drinking alcohol, smoking cigarette, plaque index and calculus index and chewing chat has no significant association with *Streptococcus mutans* dental caries. The reason for this discrepancy might be short data collection period and small number of the study population.

## **5. CONCLUSION AND RECOMMENDATION**

Dental caries of streptococcus mutans is the common public health problem among patients attending at dental clinic of Jimma University specialized hospital. Soft drinks, oral debris and gingival index are the associated risk factor for *Streptococcus mutans* dental caries. Health education on oral hygiene, dietary habit and dental clinic visit should be given to prevent and control *Streptococcus mutans* dental caries.

## **AUTHORS' CONTRIBUTIONS**

DS-performed the laboratory activities. DS- analyzed the data. TA-wrote the manuscript. All authors read and approved the final manuscript.

## **REFERENCES**

- Nada HA, Al-Mudallal, Essam FA, Al-Jumaily, Nidhal AA, Muhimen and Abd Al-Wahid Al-Shaibany (2008). Isolation and identification of mutan's streptococci bacteria from human dental plaque samples. *Journal of Al-Nahrain University*, 11(3), 98-105
- Catalán MA, Scott-Anne K, Klein MI, Koo H, Bowen WH, Melvin JE. (2011). Elevated Incidence of Dental Caries in a Mouse Model of Cystic Fibrosis Sample. *PLoS ONE*, 6(1):e16549.
- Pannu P, Gambhir R, Sujlana A. (2013). Correlation between the salivary *Streptococcus mutans* levels and dental caries experience in adult population of Chandigarh, India. *European Journal of Dentistry*, 7(2), 191-195.
- Kt S, Kmk M, NB, Jimson S, RS. (2013). Dental Caries Vaccine – A Possible Option? *Journal of Clinical and Diagnostic Research*, 7(6), 1250-1253.
- Whiley RA. and Beighton D. (2013). "Streptococci and Oral Streptococci." Bite-Sized Tutorials. *N.p. Web*.
- Jav ND, Webed M, Chaudhry S, Butt S, Ijaz S, Asad R, Awais F, and Khan A. (2013). "Transmission of *Streptococcus mutans* from Mother to Child." Review Article. *Pakistan and Oral Dental Journal*, 32 (3).
- Smith B, Pickard HM, Kidd EAM. (1990). "Why restore teeth?" *Pickard's manual of operative dentistry*, (6th Ed).
- Zhang L, Foxman B, Drake DR, et al. (2009). Comparative whole-genome analysis of *Streptococcus mutans* isolates within and among individuals of different caries status. *Oral microbiology and immunology*, 24(3), 197-203.
- Teresa CC, Franco EF, Patrícia A, José Moacir M, Fernando ADA. (2007). Detection of *Streptococcus mutans* and *Streptococcus sobrinus* in Dental Plaque Samples from Brazilian Preschool Children by Polymerase Chain Reaction *Brazil Dental Journal*, 18(4), 329-333
- Cockerill FR, Wikler MA, Bush K, Dudley MN, Eliopoulos GM, Hardy DJ, et al. (2011). Performance Standards for Antimicrobial Susceptibility Testing Twenty-First Informational Supplement.
- Costa SM, Vasconcelos M, Haddad JPA, Abreu MHN. (2012). the severity of dental caries in adults aged 35 to 44 years residing in the metropolitan area of a large city in Brazil: a cross-sectional study. *BMC Oral Health*, 12, 25.

- Liu L, Zhang Y, Wu W, Cheng M, Li Y, Cheng R. (2013). Prevalence and Correlates of Dental Caries in an Elderly Population in Northeast China. *PLoS ONE*, 8(11).
- Perera PJ, Abeyweera NT, Fernando MP, Warnakulasuriya TD, Ranathunga N.(2012). Prevalence of dental caries among a cohort of preschool children living in Gampaha district, Sri Lanka: A descriptive cross sectional study. *BMC Oral Health*, 12, 49.
- Al-Darwish M, El Ansari W, Bener A. (2014). Prevalence of dental caries among 12–14year old children in Qatar. *The Saudi Dental Journal*, 26(3), 115-125.
- Ayele FA, Taye BW, Ayele TA, Gelaye KA. (2013). Predictors of Dental caries among children 7–14 years old in Northwest Ethiopia: a community based cross-sectional study. *BMC Oral Health*, 13:7.
- Sukhabogi J, Parthasarathi P, Anjum S, Shekar B, Padma C, Rani A. (2014). Dental Fluorosis and Dental Caries Prevalence among 12 and 15-Year-Old School Children in Nalgonda District, Andhra Pradesh, India. *Annals of Medical and Health Sciences Research*, 4(Suppl 3).
- Prasai Dixit L, Shakya A, Shrestha M, Shrestha A. (2013). Dental caries prevalence, oral health knowledge and practice among indigenous Chepang school children of Nepal. *BMC Oral Health*, 13, 20.
- Mafuvadze BT, Mahachi L, Mafuvadze B. (2013). Dental caries and oral health practice among 12 year old school children from low socio-economic status background in Zimbabwe. *The Pan African Medical Journal*, 14, 164.
- T S. Kumar, BS, Datta M, V.T. H, Nisha VA. (2013). Prevalence, Severity and Associated Factors of Dental Caries in 3-6 Year Old Children. *Journal of Clinical and Diagnostic Research*, 7(8), 1789-1792.
- Nobile CG, Fortunato L, Bianco A, Pileggi C, Pavia M. (2014). Pattern and severity of early childhood caries in Southern Italy: a preschool-based cross-sectional study. *BMC Public Health*, 14-206.

**Citation:** Demissew Shenkute & Tsegahun Asfaw "*Streptococcus mutans* Dental Carries among Patients Attending Jimma University Specialized Hospital, Ethiopia" *International Journal Of Research Studies In Biosciences (Ijrsb)*, Vol. 7, no. 2, pp. 49-55, 2019. <http://Dx.Doi.org/10.20431/2349-0365.0702005>

**Copyright:** © 2019 Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.