

## **Acute Toxicity of Effluent from Electroplating Industry to the Common Carp, *Cyprinus Carpio* Linn.**

**V.J.Florence Borgia, \*A.J.Thatheyus**

PG and Research Department of Zoology,  
J. A. College for Women, Periyakulam, Tamilnadu, India.  
*jemijefferin@gmail.com*

\*PG and Research Department of Zoology,  
The American College, Madurai, Tamilnadu, India.  
*jthatheyus@yahoo.com*

---

**Abstract:** *Effluent samples from electroplating industry were collected and brought to the laboratory immediately for the analysis of physico-chemical characteristics. The analyzed parameters were compared with the Central Pollution Control Board (CPCB) standards. Static acute toxicity test was performed for a period of 96 hours using different concentrations of effluent of electroplating industry with the objective of evaluating the acute toxicity to the fresh water fish, *Cyprinus carpio*. The 24, 48, 72 and 96 hr LC<sub>50</sub> values were 0.203, 0.158, 0.132, and 0.128% respectively. The need of toxicity evaluation assay for confirming the quality of effluent from the point of effective environmental safe limits and to ensure integrity of aquatic environment is stressed.*

**Keywords:** *Acute toxicity, Common Carp, *Cyprinus carpio*, Electroplating industry effluent, LC<sub>50</sub>.*

---

### **1. INTRODUCTION**

Water, a universal solvent, is an essence for life on earth and hence, discharge of various toxic chemicals and substances into water makes life difficult. Fishes are aquatic and poikilothermic animals and their existence and performance is influenced by the quality of their environment [1]. With the advent of agricultural and industrial revolution, most of the water sources are becoming contaminated [2]. Uncontrolled discharge of industrial effluents containing toxic and hazardous substances, including heavy metals [3-4], led to severe impact on ecological balance and appreciable environmental deterioration. The indiscriminate discharge of effluents from electroplating industries into the natural aquatic systems poses a serious threat to the flora and fauna including fish [5]. Heavy metals are among the pollutants which build up in the food chain and they are responsible for the adverse effects and finally death of aquatic organisms [6-7]. When hazardous substances are released into the environment, an evaluation is necessary to determine the possible impact of these substances on human health and other biota [6-7]. Physico-chemical parameters are generally used for evaluation of effluent quality. However, these parameters alone cannot give a quantitative measure of the impact of pollution. Toxicity evaluation is an important and cost effective tool in wastewater quality monitoring as it provides the complete response of test organisms to all the compounds in a cumulative way [8-9]. Therefore the ultimate aim of toxicity evaluation is to predict the acceptable levels of toxicants in the environment to the biota. The objective of this study was to evaluate the acute toxicity of the electroplating effluent to *C. carpio*. *C. carpio* is having commercial importance and is a known bioindicator in toxicological testing [10-11].

### **2. MATERIALS AND METHODS**

Healthy *Cyprinus carpio*, irrespective of sex were collected from a local fish farm and they were acclimatized to laboratory conditions in well aerated dechlorinated tap water for fifteen days in fibre tanks (150 liter capacity). The water was changed once in two days to remove the metabolic wastes. After acclimatization, those with an average weight of 25-30g were selected for the study and the feeding was suspended two days before start and throughout the experiment. The effluent from

electroplating industry was collected and transported immediately to the laboratory and stored in a refrigerator and the physico-chemical parameters of the effluent sample were estimated adopting standard methods [12].

Preliminary screening test was carried out to ascertain the range of the effluent concentrations before the actual bioassay. The definitive concentrations used for the acute toxicity test were 0.1, 0.13, 0.14, 0.15, 0.16, 0.17, 0.20, 0.21, 0.23, 0.24, 0.3, 0.4, 0.5 and 0.6% in addition to the control. Ten fish were exposed to each concentration and the acute test was conducted for 96 hours. The mortality of fish after 24, 48, 72 and 96 hours exposure was recorded and subjected to probit analysis. The median lethal concentration ( $LC_{50}$ ) values for 24, 48, 72 and 96 hours were calculated with the 95% fiducial limits.

### 3. RESULTS AND DISCUSSION

The physico-chemical characteristics of the electroplating effluent are shown in Table 1. The pH was highly acidic and the levels of dissolved solids, chromium, nickel, lead, phosphates, sodium, potassium, iron, magnesium, and calcium were very high. The results of the analysis however implicated the effluent to be unsafe and deleterious to aquatic organisms when compared with the permissible limits [13-14].

**Table 1.** Physico-chemical Parameters of the Electroplating Industrial Effluent


S.No	Parameters	*CPHEEO/CPCB permissible level	Electroplating effluent
1	pH	6.0 to 9.0	< 1
2	Turbidity (NTU)	2.5 NTU	< 1
3	Total dissolved solids (mg/l)	500	105320
4	Chlorides (mg/l)	200	291
5	Sulphates (mg/l)	200	3571
6	Fluoride (mg/l)	1	400
7	Sodium (mg/l)	–	4491.95
8	Copper (mg/l)	3.0	335
9	Cadmium (mg/l)	1.0	< 0.05
10	Chromium (mg/l)	2.0	36756
11	Lead (mg/l)	0.05	11.64
12	Nickel (mg/l)	3.0	27562
13	Potassium (mg/l)	–	289.35
14	Manganese (mg/l)	–	0.94
15	Iron (mg/l)	0.3	16.48
16	Aluminium (mg/l)	–	9.84
17	Calcium (mg/l)	70	265
18	Magnesium (mg/l)	0.4	107

\*CPHEEO-Central Public Health Environment Engineering Organization

CPCB-Central Pollution Control Board

The  $LC_{50}$  values based on probit analysis were found to be 0.203, 0.158, 0.132 and 0.128% respectively for 24, 48, 72 and 96 hours (Fig 1; Table 2). There was a decline in  $LC_{50}$  values from 24 hours to 96 hours. The data indicated a decrease in  $LC_{50}$  value with the increase in duration of exposure. Similar result was obtained in *Channa striatus* [15], in the freshwater fish, *Rasbora daniconius* [16], in *Mystus cavasius* [5] and in *Oreochromis mossambicus* [17]. The  $LC_{50}$  of effluent for 96 hours (0.128%) was 1.59 times less than the  $LC_{50}$  of 24hrs (0.203%). The mortality rate of *C. carpio* remained directly proportional to duration of exposure, concentration and toxicity factor as already observed in *C. carpio* [18].

**Fig 1: Probit mortality of *Cyprinus carpio* in relation to different concentrations of Electroplating industry effluent exposed for 24, 48, 72 and 96 hours**



**Table 2. Acute toxicity test results of electroplating industrial effluent to *Cyprinus carpio*.**

Hours	Lethal concentration values (ppm)							95% Fiducial limits of LC <sub>50</sub>		Probit Regression Equation	Chi-Square Values		
	LC5	LC10	LC16	LC50	LC84	LC90	LC99	Lower	Upper	Y= a + b x	Observed	Table	Significance
24	0.125	0.139	0.151	0.203	0.271	0.295	0.399	0.185	0.222	Y= 10.47 + 7.88 x	13.75	9.49	S
48	0.080	0.093	0.105	0.158	0.239	0.269	0.413	0.146	0.172	Y= 9.47 + 5.59 x	16.45	12.59	S
72	0.075	0.085	0.094	0.132	0.184	0.203	0.289	0.117	0.148	Y= 10.99 + 6.81 x	25.61	11.07	S
96	0.082	0.091	0.098	0.128	0.168	0.182	0.241	0.102	0.161	Y= 12.57 + 8.49 x	14.27	5.99	S

S – Significant

Many workers have conducted experiments to determine the LC<sub>50</sub> value of different industrial effluents to freshwater fish. The 96hr LC<sub>50</sub> values of 6.09, 8.35 and 8.25% of paper mill effluent were reported for *Anabas testudineus*, *Channa punctats* and *Clarias batrachus* respectively [19], 0.018% detergent effluent to *Clarias gariepinus* [7], 7.07% tannery effluent to *Labeo rohito* [20], 4% electroplating industrial effluent to *Oreochromis mossambicus* [17] and 6% electroplating industrial effluent to *Labeo rohita* [21]. This clearly indicates that freshwater fish are sensitive to several industrial effluents, and based on our results *C. carpio* is sensitive to electroplating industrial effluent.

Indiscriminate discharge of electroplating effluent would be deleterious to aquatic life even at low concentration. Hence effluent treatment measures have to be implemented properly to keep our environment clean. For the removal of toxic metals, industrial effluents can be subjected to coagulation by ferric salt [22], membrane processes [23], polyaluminium chloride, chitosan and montmorillonite [24] and flocculation [25] in addition to biosorption [26].

#### 4. CONCLUSION

*C. carpio* was sensitive to even low concentrations of the electroplating industry effluent because of excessive levels of total dissolved solids, sodium, copper, chromium and nickel. Hence the effluent has to be treated before its discharge in to the environment.

#### ACKNOWLEDGMENT

The authors profusely thank the Management of Jayaraj Annapackiam College for Women (Autonomous) Periyakulam and the University Grants Commission, New Delhi for their support.

---

## REFERENCES

- [1]. Bakde C., and Poddar A.N., Effect of steel plant effluent on acid and alkaline phosphatases of gills, liver and gonads of *Cyprinus carpio* Linn (1758), Int. J. Environ sci. (6), 1305-1316 (2011).
- [2]. Khare S., and Singh S., Histopathological lesions induced by copper sulphate and lead nitrate in the gills of fresh water fish Nandus, J.Ecotoxicol. Environ.Monit. 12, 105-111 (2002).
- [3]. Gbem T.T., Balogun J.K., Lawaland F.A., and Annune P.A., Trace metal accumulation in *Clarias gariepinus* Teugules exposed to sublethal levels of tannery effluent, Sci. Total Environ. 271, 1-9 (2001).
- [4]. Woodling J.D., Brinkman S.F., and Horn B.J., Non-uniform accumulation of cadmium and copper in kidney's of wild brown trout *Salmo trutta* populations, Arch. Environ. Contam. Toxicol. 40, 318-385 (2001).
- [5]. Palanisamy P., Sasikala G., Mallikaraj D., and Natarajan G.M., Study of behavioural response of air – breathing catfish *Mystus cavasius* (Hamilton) exposed to electroplating industrial effluent chromium, Int. J. Pharm. Sci. Res. 2(7), 1790-1792 (2011).
- [6]. Farkas A., Salanki J., and Specziar A., Relation between growth and the heavy metal concentration in organs of bream *Abramis brama* L. populating Lake Balaton, Arch. Environ. Contam. Toxicol. 43, 236-243 (2002).
- [7]. Ogundiran M.A., Fawole O.O., Adewoye S.O., and Ayandiran T.A., Toxicological impact of detergent effluent on juvenile of African catfish (*Clarias gariepinus*) (Buchell 1822), Agric. Bio. J. N. Am. 1 (3), 330-342 (2010).
- [8]. Somashekar R.K., Gurudev M.R., and Siddaramaiah S., Somatic cell abnormalities induced by dye manufacturing industry waste waters in *Allium cepa*, Cytologia 50, 129-134 (1985).
- [9]. Tisler T., and Kocan J.Z., Toxicity evaluation of water from pharmaceutical industry to aquatic organisms, Wat. Sci Tech. 39, 71-76 (1999).
- [10]. ISO 7346/1., Water quality- determination of acute lethal toxicity of substances to a freshwater fish *Brachydanio rerio* Hamilton-Buchanan (Teleostei, Cypinidae), part 1: Static method. (1984).
- [11]. USEPA., Methods for measuring the acute toxicity of effluents and receiving waters to fresh water and marine organisms, EPA-821-R-02-01. (2002).
- [12]. APHA., Standard Methods for the Examination of Water and Wastewater, 20<sup>th</sup>Edn, APHA, AWWA, WPCF, Washington, USA, (2005).
- [13]. CPHEEO (Central Public Health & Environmental Engineering Organization) the Expert Committee, Manual on Sewerage & Sewerage Treatment (Second Edition), Ministry of Urban Development, Government of India, New Delhi, (1993).
- [14]. CPCB (Central Pollution Control Board) Municipal Solid Waste Management & Handling Rules, Ministry of Environment & Forests (MOEF), Government of India, New Delhi, (2000).
- [15]. Yadav A., Neraliya S., and Gopesh A., Acute toxicity levels and ethological responses of *Channa striatus* to fertilizer industrial wastewater, J. Environ. Bio. 28, 159 (2007).
- [16]. Pathan T.S., Sonawane D.L., and Khillare Y.K., Toxicity and behavioural changes in freshwater fish *Rasbora daniconius* exposed to paper mill effluent, Botany Res. Int. 2(4), 263-266 (2009).
- [17]. Navaraj P.S., and Kumaraguru A.K., Synergism of heavy metals on the respiration of *Oreochromis mossambicus*, Annu. Res. Rev. Bio. 4 (5), 805 – 816 (2014).
- [18]. Roopadevi H., and Somashekar R.K., Assessment of the toxicity of waste water from a textile Industry to *Cyprinus Carpio*, J. Environ. Biol. 33 (2), 167-171 (2012).
- [19]. Nanda P., Panigrahi S., Nanda B., and Behera M.K., Toxicity of paper mill effluent to fishes", Environ Ecol, 18(1), 220-222 (2000).
- [20]. Walia G.K., Handa D., Kaur H., and Kalotra R., Evaluation of genotoxic potential of tannery industry effluent in a freshwater fish, *Labeo rohita* via chromosomal aberration test, Indian J Appl. Res. 3, 557-559 (2013).
- [21]. Muley D.V., Karanjkar D.M., and Maske S.V., Impact of industrial effluents on the biochemical composition of freshwater fish *Labeo rohita*, J. Environ. Bio. 28(2), 245-249 (2007).
- [22]. Lakey D., Predictive model for drinking water treatment technology design – the efficiency of arsenic removal by in-situ formed ferric-hydroxide, Periodica polytechnic. 54, 45-51 (2010).

- [23]. Bodzek M., Inorganic micropollutants removal by means of membrane processes – state of the arts, *Ecol. Chem. Eng.* 20, 633-658 (2013).
- [24]. Zeng D., Zhai Y., Zhang S., and Ding F., Study and Application of a novel Tap Water Flocculant, *J Environ. Prot.* 3, 518 – 522 (2012).
- [25]. Yang Z., Yang H., Jiang Z., Cai T., Li H., Li A., and Cheng R., Flocculation of both anionic and cationic dyes in aqueous solutions by the amphoteric grafting flocculant carboxymethyl chitosan – graft polyacrylamide, *J Hazard. Mater.* 15, 254- 255 (2013).
- [26]. Nagashetti, V., Mahadevaraju, G. K., Muralidhar T. S., Javed A., Trivedi D., and Bhusal K. P., Biosorption of Heavy Metals from Soil by *Pseudomonas aeruginosa*, *Int. J Innov. Tech. Expl. Eng.* 2, 2278 - 3075 (2013).

#### **AUTHORS' BIOGRAPHY**



**V. Jemima Florence Borgia**, completed her B.Sc Degree in Zoology at Holy Cross College, Nagercoil, Tamil Nadu, India in 1990. She attained M.Sc Degree in Zoology at St. Xavier's College, Palayamkottai, in 1992. She was awarded M.Phil Degree with the specialization in Aquaculture by Bharathidasan University, Tiruchy, in 2006. She had worked as Lecturer in St. Anne's Community College, Periyakulam from 2001-2002. She has been rendering her service as Lecturer in the Department of Zoology, Jayaraj Annapackiam College for Women, Periyakulam, since June 2002. She has got the designation of Assistant Professor in Zoology since November 2007. So far she has published seven research papers in National Journals. She has just completed University Grants Commission sponsored Minor Research Project on Immunotoxic effects of effluent from electroplating industry on the common carp, *Cyprinus carpio*.



**Dr. A. Joseph Thatheyus**, is well known internationally for his research contributions in the field of Aquatic Toxicology and Environmental Biotechnology. He has worked on the effects of heavy metals, pesticides, industrial effluents and detergents on aquatic fauna especially fish and aquatic insects.

In his work Dr. Thatheyus combines the utility of fish, plants, microbes and bioproducts towards the remediation of effluents and waste water from industries. He has also ventured into the studies concerned with the biodiversity of plankton, butterflies and birds in a conservation perspective. He is incessantly exploring several promising possibilities to develop anticancer drugs from seaweeds and nanoparticles synthesized from marine algae.